

On the mass of a Kerr-anti-de Sitter spacetime in D dimensions

Nathalie Deruelle* and Joseph Katz**

**Institut d'Astrophysique de Paris,
GReCO, FRE 2435 du CNRS,
98 bis boulevard Arago, 75014, Paris, France
and*

*Institut des Hautes Etudes Scientifiques,
35 Route de Chartres, 91440, Bures-sur-Yvette, France*

*** Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, UK
and
The Racah Institute of Physics, Safra Campus, 91904 Jerusalem, Israel*

28 October 2004

Abstract

We show how to compute the mass of a Kerr-anti-de Sitter spacetime with respect to the anti-de Sitter background in any dimension, using a superpotential which has been derived from standard Noether identities. The calculation takes no account of the source of the curvature and confirms results obtained for black holes via the first law of thermodynamics.

*

In an interesting paper Gibbons, Lu, Page and Pope [1] generalized the Kerr-(anti)-de Sitter metrics to all dimensions $D > 5$. Furthermore Gibbons, Perry and Pope [2] calculated the angular momenta $J_{(D)i}$ of these rotating spacetimes using Komar's integrals. They also calculated the mass $E_{(D)}$ of rotating *black holes* in anti-de Sitter backgrounds using the first law of thermodynamics.

deruelle@ihes.fr
jkatz@phys.huji.ac.il

Let us emphasize that $E_{(D)}$ and $J_{(D)i}$ are classical concepts on which thermodynamics is built rather than the other way round. A Kerr-anti-de Sitter spacetime with the same mass might be produced by a rotating “star” rather than a black hole. Such a body has no quantum radiation nor a Bekenstein-Hawking entropy. One should surely be able to calculate the mass in that case because the spacetime admits a timelike Killing vector.

Here we show how to calculate the mass and the angular momenta of Kerr-anti-de Sitter metrics using the covariant KBL superpotential [3] (see [4] about the uniqueness and reliability of that superpotential).

We choose to write the D -dimensional Kerr-anti-de Sitter metrics in Kerr-Schild coordinates. In odd dimensions $D = 2n + 1$ for instance, the line element reads

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = d\bar{s}^2 + \frac{2m}{U} (h_\mu dx^\mu)^2 \quad \text{with} \quad \mu = \{0, 1, \mu_i, \phi_i\} \quad (i = 1, \dots, n) \quad (1)$$

where $d\bar{s}^2$ is the metric of the anti-de Sitter background and where the coordinates μ_i are subject to the constraint $\sum_{i=1}^{i=n} \mu_i^2 = 1$. The mass parameter m is an integration constant, h_μ is some null vector (denoted k_μ in [1-2]) depending on n rotation parameters a_i and U is a scalar function. Explicit forms of those functions can be found in Section 2 of ref [1]. These coordinates are the most convenient for our calculations. The background admits a timelike Killing vector ξ and n plane rotation Killing vectors η_i associated with one parameter displacements, say τ . ξ is normalized in such a way that $\xi^\mu \delta\tau$ is an infinitesimal time translation in a locally free falling *non rotating* inertial frame in the background. Thus for the background metric in its conventional static form when $D = 2n + 1$ for instance (equations (1.11) and (1.12) of Ref [1])

$$d\bar{s}^2 = - \left(1 + \frac{y^2}{l^2} \right) dt^2 + \frac{dy^2}{1 + y^2/l^2} + y^2 d\sigma_{(D-2)}^2$$

$$\text{where} \quad d\sigma_{(D-2)}^2 = \sum_{i=1}^{i=n} (d\tilde{\mu}_i^2 + \tilde{\mu}_i^2 d\phi_i^2) \quad \text{with} \quad \sum_{i=1}^{i=n} \tilde{\mu}_i^2 = 1, \quad (2)$$

the components of the timelike Killing vector corresponding to time translations are $(1, 0, 0, 0, \dots)$. In Kerr-Schild ellipsoidal coordinates ξ has the same components. On the other hand, in Boyer-Linquist coordinates used in Section 3 of ref [1], the components are $(\xi^0 = 1, \xi^1 = 0, \xi^{\mu_i} = 0, \xi^{\phi_i} = -a_i/l^2)$.

We now turn to the calculation of $E_{(D)}$ using the KBL superpotential $\hat{J}^{[\mu\nu]}$ (where a hat means multiplication by $\sqrt{-g}$ and brackets denote antisymmetrization) which is defined as (see [5] for a recent review and references) :

$$\hat{J}^{[\mu\nu]} = -\frac{1}{8\pi} \left(D^{[\mu} \hat{\xi}^{\nu]} - \overline{D^{[\mu} \hat{\xi}^{\nu]}} + \hat{\xi}^{[\mu} k^{\nu]} \right) \quad (3)$$

with a “Newton constant” $G_D \equiv 1$. The first term in (3) is the well known Komar superpotential density of the foreground with metric $\mathbf{g}$. The second term subtracts the

Komar superpotential of the background with metric $\bar{g}$. The vector k^ν in the third term is :

$$k^\nu = g^{\nu\rho}(\Gamma_{\rho\sigma}^\sigma - \overline{\Gamma_{\rho\sigma}^\sigma}) - g^{\rho\sigma}(\Gamma_{\rho\sigma}^\nu - \overline{\Gamma_{\rho\sigma}^\nu}). \quad (4)$$

The mass $E_{(D)}$ is given by the integral of $\hat{J}^{[\mu\nu]}$ on a $(D-2)$ surface S of radius $x^1 = r \rightarrow \infty$ in a $(D-1)$ hypersurface $x^0 = t = \text{const.}$ in Kerr-Schild coordinates :

$$E_{(D)} = \int_S d^{D-2}x \hat{J}^{[01]}. \quad (5)$$

We repeat that the integrand being covariant, the integral is a scalar which may be calculated in any convenient coordinates.

Angular momenta $J_{(D)i}$ are obtained by replacing ξ by η_i in the KBL superpotential $\hat{J}^{[01]}$. Since $\eta_i^0 = \eta_i^1 = 0$ it reduces to the Komar superpotentials and :

$$J_{(D)i} = \int_S d^{D-2}x \hat{J}_i^{[01]} \quad \text{with} \quad \hat{J}_i^{[01]} = -\frac{1}{8\pi} \left(D^{[0}\hat{\eta}_i^{1]} - \overline{D^{[0}\hat{\eta}_i^{1]}} \right). \quad (6)$$

The calculation of $\hat{J}^{[01]}$ for the mass is a mechanical procedure, which is straightforward in Kerr-Schild coordinates*. Using the asymptotic form of the metric given in [1] at leading order in r we arrive at :

$$\hat{J}^{[01]} = \frac{m}{8\pi\Xi} [(D-1)W - 1]\sqrt{g_{(D-2)}} + \mathcal{O}\left(\frac{m^2}{r}\right), \quad (7)$$

where $g_{(D-2)}$ is the determinant of the metric $d\sigma_{(D-2)}^2$ in (2) with tildes dropped and

$$\Xi \equiv \prod_{i=1}^{i=n} \Xi_i, \quad W \equiv \sum_{i=1}^{i=n} \frac{\mu_i^2}{\Xi_i}, \quad \Xi_i \equiv 1 - \frac{a_i^2}{l^2}, \quad (8)$$

with $a_n = 0$ in even dimensions $D = 2n$.

As for the calculation of $\hat{J}_i^{[01]}$ for the angular momenta it yields (the background term $\overline{D^{[0}\hat{\eta}_i^{1]}}$ does not contribute)

$$\hat{J}_i^{[01]} = \frac{m}{8\pi\Xi} \frac{a_i\mu_i^2}{\Xi_i} (D-1)\sqrt{g_{(D-2)}} + \mathcal{O}\left(\frac{m^2}{r}\right). \quad (9)$$

While the “multi-cylindrical” pairs of coordinates (μ_i, ϕ_i) were convenient in [1] to find the Kerr-anti-de Sitter metric as well as in our previous calculations, the explicit

* The useful properties are that the metric coefficients do not depend on x^0 and ϕ_i ; the $g_{\mu_i\mu_j}$ coefficients do not depend on m ; the other off-diagonal components are linear in m and hence do not enter the calculation. The fact that the vector h^μ is null further simplifies the calculation since $\sqrt{-g} = \sqrt{-\bar{g}}$.

integration of (5)(7) and (6)(9) is more easily performed in spherical coordinates, say, φ_k with $(k = 1, 2, \dots, D - 2)$. The result is

$$E_{2n} = m \frac{\mathcal{V}_{2n-2}}{4\pi\Xi} \sum_{i=1}^{i=n-1} \frac{1}{\Xi_i} \quad , \quad E_{2n+1} = m \frac{\mathcal{V}_{2n-1}}{4\pi\Xi} \left(\sum_{i=1}^{i=n} \frac{1}{\Xi_i} - \frac{1}{2} \right) \quad , \quad J_{(D)i} = m \frac{\mathcal{V}_{D-2}}{4\pi\Xi} \frac{a_i}{\Xi_i} \quad (10)$$

where $\mathcal{V}_N$ is the volume of a N -sphere.

These results confirm those given in [2]. However they were obtained using standard ideas about global Noether conservation laws associated with translational and rotational invariance of a background spacetime. They depend on the asymptotic form of the metric only and hence take no account of the source of the curvature. In [2] the mass was obtained by using the first law of black hole thermodynamics given that the entropy is identified with $1/4$ of the black hole area and the temperature with its surface gravity. Here we know the mass in the first place and assuming the same temperature as in [2] we may compute the entropy of a black hole if there is one.

Acknowledgments

N.D. thanks the Cambridge Centre for Mathematical Sciences and the Yukawa Institute for Theoretical Physics in Kyoto for their hospitality and acknowledges financial support from a CNRS-Royal Society Exchange Program. J.K. is grateful for the usual hospitality and financial support of the Institute of Astronomy in Cambridge where most of his work was performed.

Note Added in Proof

Since this work was completed, Gary Gibbons, Malcolm Perry and Chris Pope showed that the mass can also be computed by means of the Ashtekar-Magnon-Das conformal boundary formula ; see version (2) of their paper [2].

References

- [1] G.W. Gibbons, H. Lu, Don N. Page and C.N. Pope, *The General Kerr-de Sitter Metrics in All Dimensions*, hep-th/0404008.
- [2] G.W. Gibbons, M.J. Perry and C.N. Pope, *The First Law of Thermodynamics for Kerr-de Sitter Metrics*, hep-th/0408217.
- [3] J. Katz, *A note on Komar's anomalous factor*, Class. Quantum Grav., **2** (1985) 423; J. Katz, J. Bičák, D. Lynden-Bell, *Relativistic conservation laws and integral constraints for large cosmological perturbations*, Phys. Rev. **D55** (1997) 5759.
- [4] B. Julia and S. Silva, *Currents and superpotentials in classical gauge theories : II. Global aspects and the extension of affine gravity*, Class. Quantum Grav., **17** (2000) 4733, gr-qc/0005127.
- [5] N. Deruelle, J. Katz and S. Ogushi, *Conserved Charges in Einstein- Gauss-Bonnet theory*, Class. Quantum Grav., **21** (2004) 1971, gr-qc/0310098.